



IN VITRO FERTILIZATION - AS AN EFFECTIVE METHOD OF INFERTILITY TREATMENT

Mustafoeva Zarnigor Muminovna
Shokirova Munisa Kaxramon qizi

3rd year student of the Higher Nursing Faculty
Samarkand State Medical Institute, Uzbekistan

Abstract:

Reproduction for more than 7 years. About three thousand married couples passed IVF program and about 35% of them have found happiness of motherhood and fatherhood. The women with tubal-peritoneal and endocrine infertility factors and men with severe astenoteratozoospermia became parents, because the using of IVF and method intracytoplasmic sperm injection allows to fertilize eggs by single sperm. The bank of donor sperm is functioning in the Center of planning for the 9 years.

Key words:

In vitro fertilization and embryo transfer, chorionic gonadotropin, functional uterus



During IVF, mature eggs are collected (retrieved) from ovaries and fertilized by sperm in a lab. Then the fertilized egg (embryo) or eggs (embryos) are transferred to a uterus. One full cycle of IVF takes about three weeks. Sometimes these steps are split into different parts and the process can take longer.

The in vitro fertilization process breaks down into three essential components: induction of ovulation, fertilization of the oocyte, and development of embryos that are transferred into the uterus. Problems may arise resulting in failure at any one of these junctions. In 1984, the World Congress on In Vitro Fertilization was held, looking at 9,641 laparoscopies yielding 1,101 clinical pregnancies, with an overall pregnancy rate of 11 percent--clearly indicating that in vitro fertilization/embryo transfer (IVF/ET) was an idea whose time had come. Ovulation induction is monitored by both the use of ultrasound and daily estradiol levels, ultrasound indicating the number of oocytes that will be available for capture, and estradiol indicating in an indirect way the quality of those oocytes. It is a major aim in each patient to obtain at least four embryos, since this optimizes success rates. Ovulation induction at Yale is carried out with a high-dose human menopausal gonadotropin (HMG)/human chorionic gonadotropin (HCG) regimen. This regimen has insured us a success rate of 17 percent clinical pregnancies per laparoscopy. In the future, modification will occur in the process with cryopreservation of oocytes and embryos, and gamete manipulation. The modifications will be effected primarily to increase pregnancy rates. Research will continue mainly to delineate better biochemical markers for oocyte quality, but also to further explain the mystery of implantation.

Sometimes, IVF is offered as a primary treatment for infertility in women over age 40. IVF can also be done if you have certain health conditions. For example, IVF may be an option if you or your partner has:

- Fallopian tube damage or blockage. Fallopian tube damage or blockage makes it difficult for an egg to be fertilized or for an embryo to travel to the uterus.
- Ovulation disorders. If ovulation is infrequent or absent, fewer eggs are available for fertilization.
- Endometriosis. Endometriosis occurs when the uterine tissue implants and grows outside of the uterus – often affecting the function of the ovaries, uterus and fallopian tubes.
- Uterine fibroids. Fibroids are benign tumors in the wall of the uterus and are common in women in their 30s and 40s. Fibroids can interfere with implantation of the fertilized egg.
- Previous tubal sterilization or removal. If you've had tubal ligation – a type of sterilization in which your fallopian tubes are cut or blocked to permanently prevent pregnancy – and want to conceive, IVF may be an alternative to tubal ligation reversal.
- Impaired sperm production or function. Below-average sperm concentration, weak movement of sperm (poor mobility), or abnormalities in sperm size and shape can make it difficult for sperm to fertilize an egg. If semen abnormalities are found, your partner might need to see a specialist to determine if there are correctable problems or underlying health concerns.
- Unexplained infertility. Unexplained infertility means no cause of infertility has been found despite evaluation for common causes.
- A genetic disorder. If you or your partner is at risk of passing on a genetic disorder to your child, you may be candidates for preimplantation genetic testing – a procedure that involves IVF. After the eggs are harvested and fertilized, they're screened for certain genetic problems, although not all genetic problems can be found. Embryos that don't contain identified problems can be transferred to the uterus.
- Fertility preservation for cancer or other health conditions. If you're about to start cancer treatment – such as radiation or chemotherapy – that could harm your fertility, IVF for fertility preservation may be an option. Women can have eggs harvested from their ovaries and frozen in an unfertilized state for later use. Or the eggs can be fertilized and frozen as embryos for future use.

Women who don't have a functional uterus or for whom pregnancy poses a serious health risk might choose IVF using another person to carry the pregnancy (gestational carrier). In this case, the woman's eggs are fertilized with sperm, but the resulting embryos are placed in the gestational carrier's uterus.

Literature:

1. Guide to Clinical Embryology / Ed. V.S.Korsak. - M.: Medical book. - 2011
2. Register of ART. RARCH report. - 2012
3. Secrets of reproductive medicine / T. Peter, K. Chan, M. Goldstein, Z. Rosenvox; translation from English - M.: MEDpress-inform, 2006
4. In vitro fertilization and its new directions in the treatment of female and male infertility / Ed. V.I.Kulakova, B.V. Leonova. - 2nd ed. - M.: MIA, 2004